

ISSN 2310-6697

toiser—open transactions on independent scientific-engineering research

FUNKTECHNIKPLUS # JOURNAL

Théorie—Expérimentation—Métrologie—Logiciel—Applications

ISSUE 4 – SATURDAY 31 MAY 2014 – YEAR 1

REPRINT

Spherical Beltrami Fields in Chiral Media: Reciprocity and General Theorems

Nikolaos M. Berketis



*This small European Journal is
In the Defense of Honesty in Science and Ethics in Engineering*

Publisher – otoiser—open transactions on independent scientific engineering research, www.otoiser.org—info@otoiser.org, Hauptstraße 52, 2831 Scheiblingkirchen, Austria

Language – We declare the origins of the Journal by using in the cover page English, German and French, as well as, a Hellenic vignette. However, since we recognize the dominance of US English in the technical literature, we adopted it as the Journal's language, although it is not our native language.

Focus – We consider Radio–FUNK, which still creates a vivid impression of the untouchable, and its Technology–TECHNIK, from an Advanced–PLUS point of view, Plus–PLUS Telecommunications Engineering, Electrical Engineering and Computer Science, that is, we dynamically focus at any related scientific-engineering research regarding Théorie, Expérimentation, Métrologie, Logiciel, ou Applications.

Scope – We emphasize this scope broadness by extending the title of the Journal with a Doppelkreuz-Zeichen # which we use as a placeholder for substitution of our Editorial Team disciplines: # Telecommunications etc. as above, or # High Voltage, # Software Engineering, # Simulation etc. as below.

Frequency – We publish 3 issues per year: on 31st of January, on 31st of May, and on 30th of September, as well as, an extra issue every 3 papers and a volume every 2 years.

Editions – We increase the edition number of an issue only when is needed to reform one or more of its papers—thus to increase their version numbers—but we keep unchanged its 1st edition date shown on its front page and we number its pages sequentially from 1. We count the editions of *About* separately.

Format – We use a fixed-space font, hyphenation, justification, unfixed word spacing, and the uncommon for Journals A5 (half A4) page size to achieve WYSIWYG printing and clear reading of 2 to 4 side-by-side pages on wide-screen displays

Printing-on-Demand – We can email gratis PDFs at 300-4000 dpi in booklet page scaling of either brochure or book type.

Copyright – We publish under a Creative Commons Attribution, CC-BY 3.0 Unported or CC-BY 4.0 International, License only.

Please download the latest *About* edition from
about.ftpj.otoiser.org

Editorial Team

Electrical Engineering

High Voltage Engineering # Insulating Materials
Professor Michael Danikas, mdanikas@ee.duth.gr
EECE, Democritus University of Thrace, Greece

Electrical Machines # Renewable Energies # Electric Vehicles
Assistant Professor Athanasios Karlis, akarlis@ee.duth.gr
EECE, Democritus University of Thrace, Greece

Computer Science

Computer Engineering # Software Engineering # Cyber Security
Professor Vasilis Katos, vkatos@bournemouth.ac.uk
Head of Computer and Informatics Dept, Bournemouth Univ, UK

Internet Engineering # Learning Management Systems
Lecturer Sotirios Kontogiannis, skontog@gmail.com
Business Administration Dept, TEI, Western Macedonia, Greece

Hypercomputation # Fuzzy Computation # Digital Typography
Dr. Apostolos Syropoulos, asyropoulos@yahoo.com
BSc-Physics, MSc-Computer Science, PhD-Computer Science
Independent Researcher, Xanthi, Greece

Telecommunications Engineering

Applied EM Electromagnetics # Applied Mathematics
Dr. Nikolaos Berketis, nberketis@gmail.com
BSc-Mathematics, MSc-Applied Maths, PhD-Applied Mathematics
Independent Researcher, Athens, Greece

Antennas # Metrology # EM Software # Simulation # Virtual Labs

Applied EM # Education # FLOSS # Amateur Radio # Electronics
Dr. Nikolitsa Yannopoulou, yin@arg.op4.eu *
Diploma Eng-EE, MEng-Telecom-EECE, PhD-Eng-Antennas-EECE
Independent Researcher, Scheiblingkirchen, Austria

Dr. Petros Zimourtopoulos, pez@arg.op4.eu *
BSc-Physics, MSc-Radio-Electronics, PhD-Antennas-EE
Independent Researcher, Scheiblingkirchen, Austria

* Copy & Layout Editing – Proof Reading – Issue & Web Management

Technical Support

Konstantinos Kondylis, kkondylis@gmail.com
Diploma Eng-EECE, MEng-EECE, Doha, Qatar

Christos Koutsos, ckoutsos@gmail.com
Diploma Eng-EECE, MEng-EECE, Bratislava, Slovakia

Information for Authors

This is a small, but independent, low profile Journal, in which we are all—Authors, Reviewers, Readers, and Editors—free at last to be Peers in Knowledge, without suffering from Journal roles or positions, Professional–Amateur–Academic statuses, or established "impact factorizations", under the following guiding principles:

Authors – We know what Work means, we respect the Work of the Independent Researcher in Science and Engineering and we want to exhibit his Work. Thus, we decided to found this Free and Open Access Journal in which to publish this Work. Furthermore, as we care indeed for the Work of the technical author—especially a young or a beginner one—we strongly support the publication of his Work, as follows:

- 0 We do not demand from the author to transfer his own copyright to us. Instead, we only consider papers resulting from original research work only, and only if the author can assure us that he owns the copyright of his own paper as well as that he submits to the Journal either an original copy or a revised version of his own paper, for possible publication after review—or even for immediate republication, if this paper has already been published after review—but, in any case under a Creative Commons Attribution, CC-BY 3 Unported or CC-BY 4 International, License, only.
- 1 We encourage the author to submit his own paper written just in Basic English plus Technical Terminology.
- 2 We encourage the author even to select a pen name, which may drop it at any time to reveal his identity.
- 3 We encourage the author to submit an accepted for publication paper, which he was forced to decline that publication because it would be based on a review with unacceptable evaluation or derogatory comments.
- 4 We encourage the author to submit any paper that was rejected after a poor, impotent, inadequate, unreasonable, irresponsible, incompetent, or "just ticking" review.
- 5 We encourage the author to submit an unreviewed paper of his own that he uploaded on some Open Access repository.
- 6 We encourage the author to upload his published papers in our Journal to a truly Free Open repository like viXra.org.
- 7 We provide the author with a decent, express, peer review process, of up to just 4 weeks, by at least 2, either anonymous or onymous, reviewers.

- 8 We provide the author with the option to choose from 2 review processes: the traditional, anonymous, close one, as well as, a contemporary, onymous, open review in our private mailing list for Peer Discussion.
- 9 Under the Clause 0 : We immediately accept for publication a research paper directly resulting from a Project Report, or a Diploma-, Master-, or PhD-thesis, which already the author has successfully defended before a committee of experts, as long as he can mention 2 members of this committee who approved his Work.
- 10 Under the Clause 0 : We immediately accept for publication any paper which is not Openly Accessible on the Internet.
- 11 We immediately publish online a paper, as soon as it is accepted for publication in the Journal.
- 12 We quickly publish an extra issue—that is in excess of the 3 issues we publish a year—as soon as the review process of 3 papers is completed.

Reviewers – Every peer may voluntarily become a reviewer of the Journal in his skillfulness for as long as he wishes. In addition, each author of the Journal must review one paper in his expertness for each one of his published papers.

Readers – Every reader is a potential post-reviewer: we welcome comments and post-reviews in our private mailing list for Peer Discussion.

Editors – Every editor owns a PhD degree—to objectively prove that he really has the working experience of passing through the dominant publishing system. An editor pre-reviews a paper in order to check its compliance to our guiding principles and to select the appropriate reviewers of it. We can accept for consideration papers only in the expertise areas currently shown in the Editorial Team page, above. However, since we are very willing to amplify and extend the Scope of the Journal, we welcome the volunteer expert, in any related subject, who wants to join the Editorial Team as long as he unreservedly accepts our guiding principles.

Electronic Publishing

We regularly use the Free Libre Open Source Software Libre Office with the Free Liberation Mono font and the Freewares PDFCreator and PDF-Xchange Viewer. We also use, with some basic html code of ours: the Free Open Source Software Open Journal System OJS by the Public Knowledge Project PKP installed in our website, and the Free Open Digital Library of

About

Internet Archive website, where we upload Issues, Paper reprints, *About* documents, and Volumes, in both portrait and landscape orientations, for exceptionally clear online reading with the Free Open Source BookReader.

Submissions

We can consider only papers submitted in a format which is fully compatible with LibreOffice—preferably in odt format.

Legal Notice – It is taken for granted that the submitter–correspondent author accepts, without any reservation, the totality of our publication conditions as they are analytically detailed here, in this *About*, as well as, that he also carries, in the case of a paper by multiple authors, the independent will of each one of his coauthors to unreservedly accept all the aforementioned conditions for their paper.

Internet Addresses

Internet Publishing Website : ftpj.otoiser.org

Internet Archive Digital Library : archive.org/details/@ftpj

Printing-on-Demand : pod@ftpj.otoiser.org

Principal Contact : principal-contact@ftpj.otoiser.org

Technical Support : technical-support@ftpj.otoiser.org

Editorial Team–Technical Support List : etts.ftpj.otoiser.org

Peer Discussion List : peers.ftpj.otoiser.org

Submissions : sub@ftpj.otoiser.org

Sample Paper Template : template.ftpj.otoiser.org

ARG NfP AoI

Antennas Research Group

Not-for-Profit Association of Individuals *

arg.op4.eu – arg@op4.eu

Hauptstraße 52, 2831 Scheiblingkirchen, Austria

* The Constitution of Greece, Article: 12(3) 2008:
www.hellenicparliament.gr/en/Vouli-ton-Ellinon/To-Politevma

* The Hellenic Supreme Court of Civil and Penal Law:
www.areiospagos.gr/en/ – Court Rulings:Civil|A1|511|2008

This document is licensed under a Creative Commons Attribution 4.0 International License – <https://creativecommons.org/licenses/by/4.0/>

Spherical Beltrami Fields in Chiral Media: Reciprocity and General Theorems

Nikolaos M. Berketis *

Independent Researcher, Athens, Greece

Abstract

The Beltrami fields are extremely important for the description of time-harmonic electromagnetic fields in chiral media. This paper introduces the spherical Beltrami fields in chiral media, extending the corresponding known results for plane electromagnetic waves. Two key theorems are given for the scattering of spherical electromagnetic waves in chiral environment from a perfect conductor: the Reciprocity and the General Scattering theorems.

Keywords

Chiral media, spherical Beltrami fields, reciprocity relation, general scattering theorem

Introduction

The name "Beltrami field" comes from the fact that these fields \mathbf{A} satisfy the equation type $\nabla \times \mathbf{A} = \alpha \mathbf{A}$, where $\alpha \neq 0$ is a reciprocal length [1]. The word chiral regards the concept of "chirality", derived from the Greek word hand. In geometry, a figure is chiral if it cannot be mapped to its mirror image by rotations and translations alone. Typical examples are human hands, snail shells, spirals and spirals in general.

The discoveries of natural optical activity in special

materials have been known since the beginning of last century. Though optical activity has been considered in optics and in quantum mechanics for many years, its analysis within the framework of classical electromagnetic field theory arose much later. Recently, there has been a considerable interest in the study of scattering and diffraction by chiral medium. Chiral media are isotropic birefringent substances that respond to either electric or magnetic excitation with both electric and magnetic polarizations. Such media have been known since the end of the

nineteenth century (e.g. the study of chirality by Pasteur) and find a wide range of applications from many sciences [1].

In general, the electromagnetic fields inside the chiral medium are governed by Maxwell equations together with Drude-Born-Fedorov equations in which the electric and magnetic fields are coupled [1]. The chiral medium is characterized by the electric permittivity ϵ , magnetic permeability μ and chirality measure β .

Scattering problems with incident waves have been studied: for spherical acoustic waves in the works [2]-[4], as well in the works [2], [5] for spherical electromagnetic waves in non-chiral electromagnetic waves and finally for spherical electromagnetic waves in chiral media in the works [6], [7].

In this work, we derive reciprocity relation and general scattering theorem, from spherical electromagnetic waves emanating from point sources in chiral media and scattered by a perfect conductor. Similar theorems are proved, analogously, for a chiral dielectric [6], [7].

As it is well known [1], in a homogeneous isotropic chiral medium the electromagnetic field is composed of left-circularly polarized (LCP)

and right-circularly polarized (RCP) components which are propagated independently and with different phase speeds. When either a LCP or a RCP or a linear combination of LCP and RCP electromagnetic waves is incident upon a chiral scatterer then the scattered field is composed of both LCP and RCP components. So, using Bohren decomposition [1], [6], [7], the electromagnetic waves are expressed in terms of LCP and RCP Beltrami fields. The use of the Beltrami fields for problems of scattering in chiral material leads to a more simplified relationship.

This is due to the fact that the differential equations which satisfy Beltrami fields is of first order while the electric and magnetic field satisfy the Helmholtz modified equation which is of second-order. Moreover, the Beltrami equations and the conditions of radiation for the Beltrami fields are in effect separately for each one of them but on the scatterer surface, when applying the boundary conditions, both Beltrami fields are present. This fact makes easier to use these equations mainly in scattering problems where the behavior of the scattered field away from the scatterer is studied.

In the second section, we formulate the problem for

spherical electromagnetic wave in a chiral medium. In the third section we formulate the problem of electromagnetic wave scattering from a perfect conductor as a function of the Beltrami fields using the Bohren transformation. The spherical Beltrami fields are defined in the fourth section. Finally, Reciprocity relation and the General Scattering theorem are given in fifth section.

Problem Formulation

Let Ω^- be a bounded and closed subset of \mathbb{R}^3 having a C^2 -boundary S , i.e. $\partial\Omega^- = S$. The set Ω^- will be referred to as the scatterer. The exterior $\Omega^+ = \mathbb{R}^3/\Omega^-$ of the scatterer is an infinite isotropic homogeneous chiral medium with electric permittivity ε , magnetic permeability μ and chirality measure β .

The scatterer is filled with a isotropic chiral medium with corresponding physical parameters ε^-, μ^- and β^- . All the physical parameters are assumed to be real positive constants.

Let $(\mathbf{E}_0^{inc}, \mathbf{H}_0^{inc})$ be a time-Harmonic spherical electromagnetic wave, due a point source located at a point with position vector \mathbf{r}_0 with

respect to an origin O in the vicinity of Ω^- .

This wave is incident upon the scatterer Ω^- and let $(\mathbf{E}_0^{sc}, \mathbf{H}_0^{sc})$ be the corresponding scattered field. Then the total electromagnetic field $(\mathbf{E}_0^t, \mathbf{H}_0^t)$ in Ω^+ is given by

$$\begin{cases} \mathbf{E}_0^t(\mathbf{r}) = \mathbf{E}_0^{inc}(\mathbf{r}) + \mathbf{E}_0^{sc}(\mathbf{r}) \\ \mathbf{H}_0^t(\mathbf{r}) = \mathbf{H}_0^{inc}(\mathbf{r}) + \mathbf{H}_0^{sc}(\mathbf{r}) \end{cases} \quad (1)$$

We assume that the scattered field satisfies the Silver-Müller radiation condition

$$\frac{1}{\eta} \mathbf{E}_0^{sc}(\mathbf{r}) + \hat{\mathbf{r}} \times \mathbf{H}_0^{sc}(\mathbf{r}) = o(r^{-1}), \quad (2)$$

$$r \rightarrow \infty$$

uniformly in all directions, $\hat{\mathbf{r}} \in S^2$, where S^2 is the unit sphere in \mathbb{R}^3 , $r = |\mathbf{r}|$, $\hat{\mathbf{r}} = \mathbf{r}/r$ and $\eta = (\mu/\varepsilon)^{1/2}$ is the intrinsic impedance of the chiral medium in Ω^+ . It is known [8], that (2) can be replaced by

$$\hat{\mathbf{r}} \times \mathbf{E}_0^{sc}(\mathbf{r}) - \eta \mathbf{H}_0^{sc}(\mathbf{r}) = o(r^{-1}), \quad (3)$$

$$r \rightarrow \infty$$

uniformly in all directions, $\hat{\mathbf{r}} \in S^2$. The total electric field \mathbf{E}^{tot} satisfies the boundary condition:

$$\hat{\mathbf{n}} \times \mathbf{E}^t(\mathbf{r}) = \mathbf{0}, \quad \mathbf{r} \in S \quad (4)$$

where \hat{n} is the unit vector perpendicular to the outer surface S . In view of the Drude-Born-Fedorov constitutive relations [1], the total exterior electromagnetic field satisfies in the source-free region Ω^+ the modified Maxwell equations

$$\nabla \times \mathbf{E}_{r_0}^t(\mathbf{r}) = \beta \gamma^2 \mathbf{E}_{r_0}^t(\mathbf{r}) + i\omega\mu \left(\frac{\gamma}{k}\right)^2 \mathbf{H}_{r_0}^t(\mathbf{r}) \quad (5)$$

$$\nabla \times \mathbf{H}_{r_0}^t(\mathbf{r}) = \beta \gamma^2 \mathbf{H}_{r_0}^t(\mathbf{r}) - i\omega\varepsilon \left(\frac{\gamma}{k}\right)^2 \mathbf{E}_{r_0}^t(\mathbf{r}) \quad (6)$$

where

$$k^2 = \omega^2 \varepsilon \mu, \quad \gamma^2 = \frac{k^2}{1 - k^2 \beta^2} \quad (7)$$

with $|\beta k| < 1$, [1].

We note that (in contrast to the non-chiral case) k is not a wave number, but just a parameter without physical significance which has dimensions of inverse length and $\omega > 0$ is the angular frequency. The system of relations (5), (6) can be written as

$$\begin{bmatrix} \nabla \times \mathbf{E}_{r_0}^t \\ \nabla \times \mathbf{H}_{r_0}^t \end{bmatrix} = \left(\frac{\gamma}{k}\right)^2 \begin{bmatrix} k^2 \beta & i\omega\mu \\ -i\omega\varepsilon & k^2 \beta \end{bmatrix} \begin{bmatrix} \mathbf{E}_{r_0}^t \\ \mathbf{H}_{r_0}^t \end{bmatrix} \quad (8)$$

Beltrami Fields

As it is well known [1], [10], in chiral media LCP and RCP waves can both propagate

independently and with different phase speeds. So, we consider the Bohren [9], decomposition of \mathbf{E}_{r_0} and \mathbf{H}_{r_0} into suitable Beltrami fields \mathbf{Q}_{L,r_0} and \mathbf{Q}_{R,r_0} , as follows

$$\mathbf{E}_{r_0}^t = \mathbf{Q}_{L,r_0} + \mathbf{Q}_{R,r_0} \quad (9)$$

$$\mathbf{H}_{r_0}^t = \frac{1}{i\eta} (\mathbf{Q}_{L,r_0} - \mathbf{Q}_{R,r_0}) \quad (10)$$

where $\eta = (\mu/\varepsilon)^{1/2}$ is the impedance of the material.

Because the transformation Bohren the relation (8) is written:

$$\begin{bmatrix} \nabla \times \mathbf{Q}_{L,r_0} \\ \nabla \times \mathbf{Q}_{R,r_0} \end{bmatrix} = \begin{bmatrix} \gamma_L & 0 \\ 0 & -\gamma_R \end{bmatrix} \begin{bmatrix} \mathbf{Q}_{L,r_0} \\ \mathbf{Q}_{R,r_0} \end{bmatrix} \quad (11)$$

or equivalent

$$\begin{cases} \nabla \times \mathbf{Q}_{L,r_0} = \gamma_L \mathbf{Q}_{L,r_0} \\ \nabla \times \mathbf{Q}_{R,r_0} = -\gamma_R \mathbf{Q}_{R,r_0} \end{cases} \quad (12)$$

where γ_L, γ_R are wave numbers for Beltrami fields $\mathbf{Q}_{L,r_0}, \mathbf{Q}_{R,r_0}$ and are given by,

$$\gamma_L = \frac{k}{1 - k\beta}, \quad \gamma_R = \frac{k}{1 + k\beta} \quad (13)$$

respectively. Applying the transformation Bohren we split the field $\mathbf{E}_{r_0}^t, \mathbf{H}_{r_0}^t$ into the Beltrami fields $\mathbf{Q}_L, \mathbf{Q}_R$, so that the problem (2)-(6) becomes equivalent to the following problem: We are

looking for \mathbf{Q}_L , \mathbf{Q}_R in Ω^+ such that

$$\nabla \times \mathbf{Q}_L(\mathbf{r}) = \gamma_L \mathbf{Q}_L(\mathbf{r}), \quad \mathbf{r} \in \Omega^+ \quad (14)$$

$$\nabla \times \mathbf{Q}_R(\mathbf{r}) = -\gamma_R \mathbf{Q}_R(\mathbf{r}), \quad \mathbf{r} \in \Omega^+ \quad (15)$$

$$\hat{\mathbf{n}} \times \mathbf{Q}_L(\mathbf{r}) = i\sqrt{\frac{\mu}{\varepsilon}} \hat{\mathbf{n}} \times \mathbf{Q}_R(\mathbf{r}), \quad \mathbf{r} \in S \quad (16)$$

$$\hat{\mathbf{r}} \times \mathbf{Q}_L(\mathbf{r}) + i\mathbf{Q}_L(\mathbf{r}) = o\left(\frac{1}{r}\right), \quad r \rightarrow \infty \quad (17)$$

$$\hat{\mathbf{r}} \times \mathbf{Q}_R(\mathbf{r}) - i\mathbf{Q}_R(\mathbf{r}) = o\left(\frac{1}{r}\right), \quad r \rightarrow \infty \quad (18)$$

Spherical Beltrami Fields

So, we consider the Bohren decomposition of $\mathbf{E}_{r_0}^{inc}$ and $\mathbf{H}_{r_0}^{inc}$ into suitable incident spherical Beltrami fields \mathbf{Q}_{L,r_0}^{inc} and \mathbf{Q}_{R,r_0}^{inc} which have the form [6], [7],

$$\mathbf{Q}_{L,r_0}^{inc}(\mathbf{r} | \hat{\mathbf{p}}_L) = A_L \tilde{\mathbf{B}}_L(\mathbf{r}, \mathbf{r}_0) \cdot \hat{\mathbf{p}}_L \quad (19)$$

$$\mathbf{Q}_{R,r_0}^{inc}(\mathbf{r} | \hat{\mathbf{p}}_R) = A_R \tilde{\mathbf{B}}_R(\mathbf{r}, \mathbf{r}_0) \cdot \hat{\mathbf{p}}_R \quad (20)$$

where

$$\tilde{\mathbf{B}}_L(\mathbf{r}, \mathbf{r}_0) = \frac{iky_L}{8\pi y^2} (\gamma_L \tilde{\mathbf{I}} + \frac{1}{\gamma_L} \nabla \nabla + \nabla \times \tilde{\mathbf{I}}) h(\gamma_L |\mathbf{r} - \mathbf{r}_0|) \quad (21)$$

$$\tilde{\mathbf{B}}_R(\mathbf{r}, \mathbf{r}_0) = \frac{iky_R}{8\pi y^2} (\gamma_R \tilde{\mathbf{I}} + \frac{1}{\gamma_R} \nabla \nabla - \nabla \times \tilde{\mathbf{I}}) h(\gamma_R |\mathbf{r} - \mathbf{r}_0|) \quad (22)$$

and $h(x) = e^{ix}/(ix)$ is the zeroth order spherical Hankel function of the first kind and $\tilde{\mathbf{I}} = \hat{\mathbf{x}}\hat{\mathbf{x}} + \hat{\mathbf{y}}\hat{\mathbf{y}} + \hat{\mathbf{z}}\hat{\mathbf{z}}$ is the identity dyadic. We recall that $\tilde{\mathbf{B}}_L$ and $\tilde{\mathbf{B}}_R$ are the fundamental Green dyadics, for the Beltrami fields [1], [11]. The constant unit vectors $\hat{\mathbf{p}}_L$ and $\hat{\mathbf{p}}_R$ are assumed to satisfy the relations:

$$\begin{aligned} \hat{\mathbf{r}}_0 \cdot \hat{\mathbf{p}}_L &= \hat{\mathbf{r}}_0 \cdot \hat{\mathbf{p}}_R = 0 \\ \hat{\mathbf{r}}_0 \times \hat{\mathbf{p}}_L &= i\hat{\mathbf{p}}_L \\ \hat{\mathbf{r}}_0 \times \hat{\mathbf{p}}_R &= -i\hat{\mathbf{p}}_R \end{aligned} \quad (23)$$

The constants A_L and A_R are evaluated so that as the location of the point source goes to infinity along the ray in the direction $\hat{\mathbf{r}}_0$, the spherical fields degenerate into plane LCP and RCP Beltrami fields propagated in a direction from \mathbf{r}_0 towards 0 , with polarizations $\hat{\mathbf{p}}_L$ and $\hat{\mathbf{p}}_R$, respectively. Using the asymptotic forms:

$$\begin{aligned} |\mathbf{r} - \mathbf{r}_0| &= r_0 - \hat{\mathbf{r}}_0 \cdot \mathbf{r} + o\left(\frac{1}{r_0}\right), \\ \frac{\mathbf{r} - \mathbf{r}_0}{|\mathbf{r} - \mathbf{r}_0|} &= -\hat{\mathbf{r}}_0 + o\left(\frac{1}{r_0}\right), \end{aligned} \quad (24)$$

$$r_0 \rightarrow \infty$$

in the relations (19) and (20) we obtain [6], [7],

$$\mathbf{Q}_{j,r_0}^{inc}(\mathbf{r} | \hat{\mathbf{p}}_j) = A_j \frac{2e^{-iy_j r_0}}{r_0} \tilde{\mathbf{K}}_j(-\hat{\mathbf{r}}_0) \cdot \hat{\mathbf{p}}_j e^{-iy_j \hat{\mathbf{r}}_0 \cdot \mathbf{r}} + O\left(\frac{1}{r_0^2}\right), \quad (25)$$

$$r_0 \rightarrow \infty$$

for $j = L, R$, where

$$\tilde{\mathbf{K}}_L(-\hat{\mathbf{r}}_0) = \frac{1}{2}(\tilde{\mathbf{I}} - \hat{\mathbf{r}}_0 \hat{\mathbf{r}}_0 - i\hat{\mathbf{r}}_0 \times \tilde{\mathbf{I}}) \quad (26)$$

$$\tilde{\mathbf{K}}_R(-\hat{\mathbf{r}}_0) = \frac{1}{2}(\tilde{\mathbf{I}} - \hat{\mathbf{r}}_0 \hat{\mathbf{r}}_0 + i\hat{\mathbf{r}}_0 \times \tilde{\mathbf{I}}) \quad (27)$$

In view of (23) the dyadics $\tilde{\mathbf{K}}_L$ and $\tilde{\mathbf{K}}_R$ satisfy the relations

$$\begin{cases} \hat{\mathbf{p}}_L = \tilde{\mathbf{K}}_L(-\hat{\mathbf{r}}_0) \cdot \hat{\mathbf{p}}_L \\ \hat{\mathbf{p}}_R = \tilde{\mathbf{K}}_R(-\hat{\mathbf{r}}_0) \cdot \hat{\mathbf{p}}_R \end{cases} \quad (28)$$

Hence, if we take

$$\begin{cases} A_L = r_0 e^{-iy_L r_0} \frac{4\pi y^2}{k y_L} \\ A_R = r_0 e^{-iy_R r_0} \frac{4\pi y^2}{k y_R} \end{cases} \quad (29)$$

then

$$\begin{aligned} \lim_{r_0 \rightarrow \infty} \mathbf{Q}_{L,r_0}^{inc}(\mathbf{r} | \hat{\mathbf{p}}_j) &= e^{-iy_L \hat{\mathbf{r}}_0 \cdot \mathbf{r}} \hat{\mathbf{p}}_j = \\ &= \mathbf{Q}_j^{inc}(\mathbf{r}; -\hat{\mathbf{r}}_0, \hat{\mathbf{p}}_j) \end{aligned} \quad (30)$$

It is convenient to write the incident spherical Beltrami fields as [6], [7],

$$\mathbf{Q}_{L,r_0}^{inc}(\mathbf{r} | \hat{\mathbf{p}}_L) = \tilde{\mathbf{r}}_L \left(\frac{h(y_L u)}{h(y_L r_0)} \right) \cdot \hat{\mathbf{p}}_L \quad (31)$$

$$\mathbf{Q}_{R,r_0}^{inc}(\mathbf{r} | \hat{\mathbf{p}}_R) = \tilde{\mathbf{r}}_R \left(\frac{h(y_R u)}{h(y_R r_0)} \right) \cdot \hat{\mathbf{p}}_R \quad (32)$$

where

$$\tilde{\mathbf{r}}_L = \frac{1}{2y_L}(y_L \tilde{\mathbf{I}} + \frac{1}{y_L} \nabla \nabla + \nabla \times \tilde{\mathbf{I}}) \quad (33)$$

$$\tilde{\mathbf{r}}_R = \frac{1}{2y_R}(y_R \tilde{\mathbf{I}} + \frac{1}{y_R} \nabla \nabla - \nabla \times \tilde{\mathbf{I}}) \quad (34)$$

and $u = |\mathbf{r} - \mathbf{r}_0|$. Using asymptotic forms (24) for $r \rightarrow \infty$ we obtain [6], [7],

$$\mathbf{Q}_{j,r_0}^{inc}(\mathbf{r} | \hat{\mathbf{p}}_j) = \mathbf{F}_{j,r_0}^{inc}(\hat{\mathbf{r}} | \hat{\mathbf{p}}_j) h(y_j r) + O\left(\frac{1}{r^2}\right) \quad (35)$$

where

$$\mathbf{F}_{j,r_0}^{inc}(\hat{\mathbf{r}} | \hat{\mathbf{p}}_j) = \frac{e^{-iy_j \hat{\mathbf{r}} \cdot \mathbf{r}_0}}{h(y_j r_0)} \tilde{\mathbf{K}}_j(\hat{\mathbf{r}}) \cdot \hat{\mathbf{p}}_j \quad (36)$$

are the far-field patterns of the point source incident Beltrami fields, which satisfy the relations

$$\hat{\mathbf{r}} \cdot \mathbf{F}_{j,r_0}^{inc}(\hat{\mathbf{r}} | \hat{\mathbf{p}}_j) = 0 \quad (37)$$

The scattered electric field $\mathbf{E}_{r_0}^{sc}$ will be dependent on the polarization $\hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R$ and will be have the decomposition

$$\begin{aligned} \mathbf{E}_{r_0}^{sc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R) &= \mathbf{Q}_{L,r_0}^{sc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R) \\ &+ \mathbf{Q}_{R,r_0}^{sc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R) \end{aligned} \quad (38)$$

where

$\mathbf{Q}_{L,r_0}^{sc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R)$ and $\mathbf{Q}_{R,r_0}^{sc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R)$ are the corresponding scattered Beltrami fields, which have the following behavior, when $r \rightarrow \infty$ [1], [10], [12]

$$\mathbf{g}_{L,r_0}(\hat{\mathbf{r}}) = \frac{iky_L}{8\pi y^2} \tilde{\mathbf{K}}_L(\hat{\mathbf{r}}) \cdot \int_S \hat{\mathbf{n}} \times (y_L \nabla \times \mathbf{E}_{r_0}^{sc}(\mathbf{r}') + y^2 \mathbf{E}_{r_0}^{sc}(\mathbf{r}')) e^{-iy_L \hat{\mathbf{r}} \cdot \mathbf{r}'} ds(\mathbf{r}') \quad (40)$$

$$\mathbf{g}_{R,r_0}(\hat{\mathbf{r}}) = \frac{iky_R}{8\pi y^2} \tilde{\mathbf{K}}_R(\hat{\mathbf{r}}) \cdot \int_S \hat{\mathbf{n}} \times (y_R \nabla \times \mathbf{E}_{r_0}^{sc}(\mathbf{r}') - y^2 \mathbf{E}_{r_0}^{sc}(\mathbf{r}')) e^{-iy_R \hat{\mathbf{r}} \cdot \mathbf{r}'} ds(\mathbf{r}') \quad (41)$$

are the LCP and RCP far-field patterns, respectively [10], and they are dependent also on the polarization $\hat{\mathbf{p}}_L$ and $\hat{\mathbf{p}}_R$. We note that the far-field patterns $\mathbf{g}_{j,r_0}(\hat{\mathbf{r}})$ satisfy the relations

$$\begin{cases} \hat{\mathbf{r}} \cdot \mathbf{g}_{j,r_0}(\hat{\mathbf{r}}) = 0 \\ \hat{\mathbf{r}} \times \mathbf{g}_{L,r_0}(\hat{\mathbf{r}}) = -i \mathbf{g}_{L,r_0}(\hat{\mathbf{r}}) \\ \hat{\mathbf{r}} \times \mathbf{g}_{R,r_0}(\hat{\mathbf{r}}) = i \mathbf{g}_{R,r_0}(\hat{\mathbf{r}}) \end{cases} \quad (42)$$

Reciprocity and General Scattering Theorems

For two vector functions \mathbf{u} and \mathbf{v} we introduce the bilinear form [10],

$$\begin{aligned} \{\mathbf{u}, \mathbf{v}\}_S &= \\ &= \int_S \hat{\mathbf{n}} \cdot (\mathbf{u} \times \nabla \times \mathbf{v} - \mathbf{v} \times \nabla \times \mathbf{u}) dS - \\ &- 2\beta y^2 \int_S \hat{\mathbf{n}} \cdot (\mathbf{u} \times \mathbf{v}) dS \end{aligned} \quad (43)$$

$$\mathbf{Q}_{j,r_0}^{sc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R) = h(y_j r) \mathbf{g}_{j,r_0}(\hat{\mathbf{r}}) + O\left(\frac{1}{r^2}\right) \quad (39)$$

The functions $\mathbf{g}_{j,r_0}(\hat{\mathbf{r}})$ given by

where S is the surface of the scatterer Ω^- and $\hat{\mathbf{n}}$ is the outward unit vector on S . We, also, consider two locations for the point source, \mathbf{a} and \mathbf{b} from which the time-harmonic incident spherical electric waves

$$\begin{aligned} \mathbf{E}_\sigma^{inc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R) &= \\ &= \mathbf{Q}_{L,\sigma}^{inc}(\mathbf{r} | \hat{\mathbf{p}}_L) + \mathbf{Q}_{R,\sigma}^{inc}(\mathbf{r} | \hat{\mathbf{p}}_R) \end{aligned} \quad (44)$$

for $\sigma = \mathbf{a}, \mathbf{b}$, emanate. \mathbf{E}_σ^{sc} and the corresponding scattered electric waves \mathbf{E}_σ^{sc} of form (38), have the Bohren decomposition of (9) in terms of LCP and RCP Beltrami fields. In particular, we have

$$\begin{cases} \mathbf{E}_\sigma^{inc} = \mathbf{Q}_{L,\sigma}^{inc} + \mathbf{Q}_{R,\sigma}^{inc} \\ \mathbf{E}_\sigma^{sc} = \mathbf{Q}_{L,\sigma}^{sc} + \mathbf{Q}_{R,\sigma}^{sc} \end{cases} \quad (45)$$

and the following properties for the fields $\mathbf{Q}_{j,\sigma}^{inc}$ and $\mathbf{Q}_{j,\sigma}^{sc}$ [6], [7]:

$$\{\bar{\mathbf{Q}}_{L,a}^{inc}(\mathbf{r}; \hat{\mathbf{p}}_L), \mathbf{Q}_{L,b}^{sc}(\mathbf{r} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R)\}_S = \frac{2Y^2}{K} \int_S \hat{\mathbf{n}} \cdot (\bar{\mathbf{Q}}_{L,a}^{inc}(\mathbf{r}; \hat{\mathbf{p}}_L) \times \mathbf{Q}_{L,b}^{sc}(\mathbf{r} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R)) ds(\mathbf{r}) \quad (46)$$

$$\{\bar{\mathbf{Q}}_{L,a}^{inc}(\mathbf{r}; \hat{\mathbf{p}}_L), \mathbf{Q}_{R,b}^{sc}(\mathbf{r} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R)\}_S = 0 \quad (47)$$

$$\{\bar{\mathbf{Q}}_{R,a}^{inc}(\mathbf{r}; \hat{\mathbf{p}}_R), \mathbf{Q}_{R,b}^{sc}(\mathbf{r} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R)\}_S = -\frac{2Y^2}{K} \int_S \hat{\mathbf{n}} \cdot (\bar{\mathbf{Q}}_{R,a}^{inc}(\mathbf{r}; \hat{\mathbf{p}}_R) \times \mathbf{Q}_{R,b}^{sc}(\mathbf{r} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R)) ds(\mathbf{r}) \quad (48)$$

$$\{\bar{\mathbf{Q}}_{R,a}^{inc}(\mathbf{r}; \hat{\mathbf{p}}_R), \mathbf{Q}_{L,b}^{sc}(\mathbf{r} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R)\}_S = 0 \quad (49)$$

Now a "reciprocity theorem" for spherical electromagnetic waves in chiral media is formulated as follows [6], [7]:

For any two incident spherical electric waves $\mathbf{E}_a^{inc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R)$ and $\mathbf{E}_b^{inc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R)$ of (44) and for any scatterer in a homogeneous isotropic chiral medium, we have:

$$\begin{aligned} & \frac{ae^{-iy_L a}}{Y_L} \mathbf{Q}_{L,b}^{sc}(\mathbf{a} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R) \cdot \hat{\mathbf{p}}_L + \\ & + \frac{ae^{-iy_R a}}{Y_R} \mathbf{Q}_{R,b}^{sc}(\mathbf{a} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R) \cdot \hat{\mathbf{p}}_R = \\ & = \frac{be^{-iy_L b}}{Y_L} \mathbf{Q}_{L,a}^{sc}(\mathbf{b} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R) \cdot \hat{\mathbf{q}}_L + \\ & + \frac{be^{-iy_R b}}{Y_R} \mathbf{Q}_{R,a}^{sc}(\mathbf{b} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R) \cdot \hat{\mathbf{q}}_R \end{aligned} \quad (50)$$

Let

$\mathbf{E}_a^{inc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R)$ and $\mathbf{E}_b^{inc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R)$ be two incident spherical waves of the form (44). We define spherical far-field pattern generators for LCP and RCP spherical Beltrami fields by

$$\begin{aligned} \mathbf{G}_{j,b}(\mathbf{a} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R) &= \frac{ae^{iy_L a}}{iy_j} \cdot \\ & \cdot \left\{ \frac{1}{4\pi} \int_{S^2} \mathbf{g}_{j,b}(\mathbf{a} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R) \cdot \overline{\tilde{\mathbf{K}}_j(\hat{\mathbf{r}})} e^{iy_A \hat{\mathbf{r}} \cdot \mathbf{a}} ds(\hat{\mathbf{r}}) - \right. \\ & \left. - \mathbf{Q}_{j,b}^{sc}(\mathbf{a} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R) \right\} \end{aligned} \quad (51)$$

for $j = L, R$.

This terminology and definition is appropriate because when both the observation point and the source goes to infinity, $\mathbf{G}_{j,b}(\mathbf{a} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R)$, reduce to the far-field patterns for an incident plane electric wave propagating in the direction $-\hat{\mathbf{b}}$ and of polarizations $\hat{\mathbf{q}}_L$ for LCP and $\hat{\mathbf{q}}_R$ for RCP fields. Far-field pattern generators in acoustic and achiral electromagnetic scattering have been defined in [13]. When the point sources are transferred at infinity, the generators $\mathbf{G}_{j,b}(\mathbf{a} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R)$ are transformed into far-field patterns scattering spherical electrical wave, namely [6], [7],

$$\lim_{a \rightarrow \infty} \mathbf{G}_{j,b}(\mathbf{a} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R) = \mathbf{g}_{j,b}(-\hat{\mathbf{a}} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R) \quad (52)$$

Using this notation, the "general scattering theorem" for spherical electric waves in chiral media is formulated as follows [6], [7]:

Let

$\mathbf{E}_a^{inc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R)$ and $\mathbf{E}_b^{inc}(\mathbf{r} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R)$ be two spherical electric waves of (44) incident upon a

scatterer in a chiral medium. Then it is valid

$$\begin{aligned} & \mathbf{G}_{L,b}(\mathbf{a} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R) \cdot \hat{\mathbf{p}}_L + \mathbf{G}_{R,b}(\mathbf{a} | \hat{\mathbf{q}}_L, \hat{\mathbf{q}}_R) \cdot \hat{\mathbf{p}}_R + \\ & + \overline{\mathbf{G}_{L,a}(\mathbf{b} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R)} \cdot \hat{\mathbf{q}}_L + \overline{\mathbf{G}_{R,a}(\mathbf{b} | \hat{\mathbf{p}}_L, \hat{\mathbf{p}}_R)} \cdot \hat{\mathbf{q}}_R = \\ & = -\frac{1}{2\pi} \left\{ \int_{S^2} \frac{1}{V_L^2} \overline{\mathbf{F}_{L,a}^{sc}(\hat{\mathbf{r}})} \cdot \mathbf{F}_{L,b}^{sc}(\hat{\mathbf{r}}) dS(\hat{\mathbf{r}}) + \right. \\ & \left. + \int_{S^2} \frac{1}{V_R^2} \overline{\mathbf{F}_{R,a}^{sc}(\hat{\mathbf{r}})} \cdot \mathbf{F}_{R,b}^{sc}(\hat{\mathbf{r}}) dS(\hat{\mathbf{r}}) \right\} \quad (53) \end{aligned}$$

References

- [1] Lakhtakia A., "Beltrami Fields in Chiral Media", World Scientific, Singapore, 1993, §1-1 (p. 3), §3-1 (p. 83), §3-2.2 (p. 110), §4-1 (p. 130), §4-3, (p. 150), §5-1, (p. 164)
- [2] Athanasiadis C., Martin P.A., Stratis I.G., "On the scattering of spherical electromagnetic waves by a penetrable chiral obstacle", 6th International Workshop on Mathematical Methods in Scattering Theory and Biomedical Engineering, Tsepelovo, Greece, 2003
- [3] Charalambopoulos A., Dassios G., Kamvyssas G., "Reciprocity theorems for point source scalar scattering", In: Dassios G., Fotiadis D., Kiriaki C., Massalas C. (eds) *Mathematical Methods in Scattering Theory and Biomedical Technology*, Longman, London 1998, pp. 12-19
- [4] Dassios G, Kamvyssas G. "The impedance scattering problem for a point source field. The small resistive sphere", *Quarterly Journal of Mechanics and Applied Mathematics*, Vol. 50, 1999, pp. 321-332
- [5] Athanasiadis C., Martin P.A., Stratis I.G., "On the Scattering of Point-generated Electromagnetic Waves by a Perfectly Conducting Sphere, and Related Near-field Inverse Problems", *Zeitschrift für Angewandte Mathematik und Mechanik*, Vol. 83, Issue 2, 2003, pp. 129-136
- [6] Athanasiadis C, Berketis N., "Scattering relations for point - source excitation in chiral media", *Mathematical Methods in the Applied sciences*, Vol. 29, 2006, pp. 27-48

- [7] Berketis M. Nikolaos, "Scattering spherical Electromagnetic waves in chiral media", PhD thesis, University of Athens – Department of Mathematics, December 2007
- [8] Colton D., Kress R., "Inverse Acoustic and Electromagnetic Scattering Theory" Springer, Berlin, 1992, §6.2, (p. 153)
- [9] Bohren C. F., "Light scattering by an optically active Sphere", Chemical Physics Letters, Vol. 29, Issue 3, 1974, pp. 458-462
- [10] Athanasiadis C., "On the far field patterns for electromagnetic scattering by a chiral obstacle in a chiral environment", Journal of Mathematical Analysis and Applications, Vol. 309, 2005, pp. 517-533
- [11] Athanasiadis C., Giotopoulos S., "Scattering theorem for dyadic chiral fields", Reports on Mathematical Physics, Vol. 53, 2004, pp. 339-350
- [12] Berketis M. Nikolaos, "Direct and Related Far-Field Inverse Scattering Problems for Spherical Electromagnetic Waves in Chiral Media", FunkTechnikPlus # Journal, Issue 2, November 2013, pp. 7-18
- [13] Athanasiadis C., Martin P.A., Spyropoylos A., Stratis I.G., "Scattering relations for point source acoustic and electromagnetic waves", Journal of Mathematical Physics Vol. 43, Issue 11, 2002, pp. 5683-5697

Previous Publication in FUNKTECHNIKPLUS # JOURNAL

"Near-field Method in Solving Inverse Scattering Problem of Spherical Electromagnetic Waves in Chiral media",
Issue 3, pp. 19-25

*** About the Author**

Nikolaos Berketis, Issue 2, p. 18

[This Page Intentionally Left Blank]

[This Page Intentionally Left Blank]

[This Page Intentionally Left Blank]

In case of any doubt,
download the genuine papers from
genuine.ftpj.otoiser.org

FRONT COVER VIGNETTE

A faded synthesis of an anthemion rooted in a meandros

The thirteen-leaf is a symbol for a life tree leaf.
"Herakles and Kerberos", ca. 530–500 BC,
by Paseas, the Kerberos Painter,
Museum of Fine Arts, Boston.

www.mfa.org/collections/object/plate-153852

The simple meandros is a symbol for eternal immortality.
"Warrior with a phiale", ca. 480–460 BC,
by Berliner Maler,
Museo Archeologico Regionale "Antonio Salinas" di Palermo.

commons.wikimedia.org/wiki/File:Warrior_MAR_Palermo_NI2134.jpg